

MATH 201: LINEAR ALGEBRA
TUTORIAL AND SUGGESTED PROBLEMS FOR WEEK 6

WEEK OF SEPTEMBER 30, 2025

WEEK 6 — PROBLEMS WITH SOLUTIONS

Problem 0.1. Fill in the blanks.

(a) The *image* of a linear transformation $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is

$$\text{Im}(T) = \{ \quad \quad \quad \mid \quad \quad \quad \}$$

(b) The *kernel* of a linear transformation $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is

$$\ker(T) = \{ \quad \quad \quad \mid \quad \quad \quad \}.$$

Solution.

(a) $\text{Im}(T) = \{T(x) : x \in \mathbb{R}^m\}$, the set of all outputs of T .

(b) $\ker(T) = \{x \in \mathbb{R}^m : T(x) = 0\}$, the set of inputs sent to the zero vector.

Problem 0.2. Suppose that $T_A(\vec{x}) = A\vec{x}$ where $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix}$.

(a) Write a set of vectors that *span* the image of T_A .

(b) Write a set of vectors that *span* the kernel of T_A .

(c) What is the *minimum* number of vectors needed to span the image of T_A ?

(d) What is the *minimum* number of vectors needed to span the kernel of T_A ?

Solution.

Write $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix}$. The columns are

$$c_1 = (1, 1)^\top, \quad c_2 = (1, 2)^\top, \quad c_3 = (1, 3)^\top, \quad c_4 = (1, 4)^\top$$

(a) $\text{Im}(T_A) = \text{span}\{c_1, c_2, c_3, c_4\} = \text{span}\{(1, 1)^\top, (1, 2)^\top\}$, since any two noncollinear columns span \mathbb{R}^2 .

(b) Solve $Ax = 0$ with $x = (x_1, x_2, x_3, x_4)^\top$. Row-reduce $\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \end{bmatrix}$,
so $x_1 = x_3 + 2x_4$, $x_2 = -2x_3 - 3x_4$. Let

$$s = x_3, \quad t = x_4.$$

Then

$$x = s(1, -2, 1, 0)^\top + t(2, -3, 0, 1)^\top.$$

Hence a spanning set for $\ker T_A$ is $\{(1, -2, 1, 0)^\top, (2, -3, 0, 1)^\top\}$.

(c) $\text{rank}(A) = 2$, so the minimum number is 2.

(d) $\text{nullity}(A) = 4 - 2 = 2$, so the minimum number is 2.

Problem 0.3. Suppose that A is a square matrix. Write 8 separate statements equivalent to the statement

“ A is invertible.”

In other words, review “Summary 3.3.10” from Bretscher 4th edition.

Solution.

Possible answers (any correct eight earn full credit):

- (1) A has an inverse matrix A^{-1} .
- (2) $\det A \neq 0$ (or equivalently, A row-reduces to I).
- (3) The columns of A are linearly independent.

- (4) The columns of A span \mathbb{R}^n .
- (5) $\text{rank}(A) = n$.
- (6) $\ker A = \{0\}$.
- (7) The only solution to $Ax = 0$ is $x = 0$.
- (8) For every $b \in \mathbb{R}^n$, the system $Ax = b$ has a unique solution.
- (9) A is a product of elementary matrices.
- (10) The linear map $x \mapsto Ax$ is one-to-one and onto.

(Write any eight; keep your phrasing consistent with what you've learned in class.)

Problem 0.4. Which of the following sets are *subspaces* of \mathbb{R}^3 ?

- (a) $W = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x + y + z = 1 \right\}$.
- (b) $W = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \mid x \leq y \leq z \right\}$.
- (c) $W = \left\{ \begin{bmatrix} x + 2y + 3z \\ 4x + 5y + 6z \\ 7x + 8y + 9z \end{bmatrix} \mid x, y, z \text{ are arbitrary constants} \right\}$.

Solution.

- (a) Not a subspace: it does not contain 0 (since $0 + 0 + 0 \neq 1$) and is not closed under scaling.
- (b) Not a subspace: not closed under scaling (multiply by -1 reverses the inequalities).
- (c) Yes, it is a subspace: it is the span of the three column vectors $(1, 4, 7)^\top$, $(2, 5, 8)^\top$, $(3, 6, 9)^\top$.

Problem 0.5. Consider the following lists of vectors. For each list, determine whether the given vectors are linear independent.

- (a) $\begin{bmatrix} 7 \\ 11 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix}$
- (b) $\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 5 \\ 0 \end{bmatrix}$.
- (c) $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 6 \\ 5 \\ 4 \end{bmatrix}$.

Solution.

- (a) Dependent (one vector is the zero vector).
- (b) Dependent: the first two are multiples; the last vector is the sum of 3 times the first, 4 times the third, and 5 times the fourth.
- (c) Try to solve:

$$a \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 5 \\ 4 \end{bmatrix}.$$

We have

$$\left[\begin{array}{ccc|c} 1 & 3 & 6 & 6 \\ 1 & 2 & 5 & 5 \\ 1 & 1 & 4 & 4 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & 3 & 3 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

This indicates that $a = 3, b = 1$ is a solution. That is, labeling the vectors \vec{v}_1, \vec{v}_2 , and \vec{v}_3 , we have $\vec{v}_3 = 3\vec{v}_1 + \vec{v}_2$.

Problem 0.6. Write a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ whose kernel is the line spanned by

the vector $\vec{v} = \begin{bmatrix} -1 \\ 1 \\ 2 \end{bmatrix}$.

Solution. Suppose that the 3×3 matrix A satisfies $A\vec{v} = 0$. Let $\vec{c}_1, \vec{c}_2, \vec{c}_3$ be the column vectors of A . Then

$$\begin{aligned} A\vec{v} &= -\vec{c}_1 + \vec{c}_2 + 2\vec{c}_3 = 0 \\ \Rightarrow \vec{c}_1 &= \vec{c}_2 + 2\vec{c}_3. \end{aligned}$$

Thus

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 2 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

is one example.

Problem 0.7. Write a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ whose kernel is the plane $x + 2y + 3z = 0$ in \mathbb{R}^3 .

Solution. The image of T should be *normal* to the plane. The normal vector of this plane is $\langle 1, 2, 3 \rangle$. To see this, note that $\langle x, y, z \rangle \cdot \langle 1, 2, 3 \rangle = x + 2y + 3z = 0$ is the plane equation given. Next, recall that the image of a linear transformation is equal to the *span* of its column vectors. Thus it suffices to construct a matrix A whose image is equal to the span of $\langle 1, 2, 3 \rangle$. For example,

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{bmatrix}.$$

Problem 0.8. Consider an $n \times p$ matrix A and a $p \times m$ matrix B such that $\ker A = \{\vec{0}\}$ and $\ker B = \{\vec{0}\}$. What is $\ker AB$?

Solution.

If $\ker A = \{0\}$ and $\ker B = \{0\}$, then A is injective on its domain and B is injective on its domain. If $ABx = 0$, then $A(Bx) = 0$, hence $Bx = 0$ (injectivity of A), hence $x = 0$ (injectivity of B). Therefore $\ker(AB) = \{0\}$.

Problem 0.9. Let A be a matrix and let $B = \text{rref}(A)$.

- (a) Is $\ker A$ necessarily equal to $\ker B$? Explain.
- (b) Is $\text{Im}A$ necessarily equal to $\text{Im}B$? Explain.

Solution.

(a) $\ker A = \ker(\text{rref}(A))$. Row operations preserve the solution set of $Ax = 0$, so the nullspace is unchanged.

(b) In general $\text{Im}A \neq \text{Im}(\text{rref}(A))$. Row operations mix rows (i.e., change the span of the rows, not the columns). Counterexample: $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ has image $\text{span}\{(1, 1)^\top\}$, but $\text{rref}(A) = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ has image $\text{span}\{(1, 0)^\top\}$.

Problem 0.10. Consider two subspaces V and W of \mathbb{R}^n .

- (a) Is $V \cup W$ a subspace of \mathbb{R}^n ?
- (b) Is $V \cap W$ a subspace of \mathbb{R}^n ?

Solution.

(a) Typically no: $V \cup W$ need not be closed under addition. Example in \mathbb{R}^2 : $V = x$ -axis, $W = y$ -axis; $V \cup W$ is not a subspace.

(b) Yes: $V \cap W$ is always a subspace (it contains 0 and is closed under addition and scaling).

Problem 0.11. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^p$ be a linear transformation. Suppose that $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m \in \mathbb{R}^n$ are linearly independent. Under what conditions (on T, m, n, p) are $T(\vec{v}_1), \dots, T(\vec{v}_m)$ linearly independent?

Solution. Claim: $T(\vec{v}_i)$ are linearly independent if and only if

$$\ker T \cap \text{span} \{\vec{v}_1, \dots, \vec{v}_m\} = \{\vec{0}\}.$$

Proof: First, we prove that if $T(\vec{v}_i)$ are *not* linearly independent then $\ker T \cap \text{span} \{\vec{v}_1, \dots, \vec{v}_m\}$ is nonempty. Suppose that $\sum_i a_i T(\vec{v}_i) = 0$. Then $T(\sum_i a_i \vec{v}_i) = 0$ by linearity. Thus $\sum_i a_i \vec{v}_i \in \ker T \cap \text{span} \{\vec{v}_1, \dots, \vec{v}_m\}$. Next we prove the converse. That is, we prove that if $\ker T \cap \text{span} \{\vec{v}_1, \dots, \vec{v}_m\} = \{\vec{0}\}$ then $T(\vec{v}_i)$ are *not* linearly independent. Suppose there exists a nonzero $w \in \ker T \cap \text{span} \{\vec{v}_1, \dots, \vec{v}_m\}$. Then $w = \sum_i a_i \vec{v}_i$ and $T(w) = 0$ so we have a nontrivial relation among $T(\vec{v}_i)$ as desired. □

Problem 0.12. Find a *basis* for the image of the matrices

(a) $\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 5 \\ 1 & 3 & 7 \end{bmatrix}$

(c) $\begin{bmatrix} 0 & 1 & 2 & 0 & 0 & 3 \\ 0 & 0 & 0 & 1 & 0 & 4 \\ 0 & 0 & 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

(d) $\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 5 & 8 \end{bmatrix}$.

Solution.

(a) Columns $\{(1, 1, 1)^\top, (1, 2, 3)^\top\}$ are linearly independent; they form a basis for the image.

(b) Row-reduce $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 5 \\ 1 & 3 & 7 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & 4 \\ 0 & 0 & 0 \end{bmatrix}$, so rank = 2. A basis for the image is $\{(1, 1, 1)^\top, (1, 2, 3)^\top\}$

(the first two pivot columns).

(c) Note that the matrix is in ref. There are *three* leading ones. We take the three pivot columns:

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}.$$

(d) In this case, there are two columns and the second is not a multiple of the first. Thus they are a basis for the image.

$$\begin{bmatrix} 1 \\ 2 \\ 3 \\ 5 \end{bmatrix}, \begin{bmatrix} 5 \\ 6 \\ 7 \\ 8 \end{bmatrix}$$

Problem 0.13. Consider an $m \times n$ matrix A and an $n \times m$ matrix B with $n \neq m$ such that $AB = I_m$. Are the columns of B linearly independent? What about the columns of A ?

Solution.

If $AB = I_m$, then B has linearly independent columns (since $ABx = 0 \Rightarrow x = 0$). Also A has linearly independent rows (equivalently, rank m). The columns of A need not be independent when $n \neq m$ (e.g., A is $m \times n$ with $m < n$ can have dependent columns).

Problem 0.14. Suppose that $\vec{v}_1, \vec{v}_2, \vec{v}_3$ in \mathbb{R}^n are linearly independent. Are the vectors $\vec{v}_1, \vec{v}_1 + \vec{v}_2, \vec{v}_1 + \vec{v}_2 + \vec{v}_3$ linearly independent?

Solution.

Yes, they are linearly independent if v_1, v_2, v_3 are independent. Suppose $av_1 + b(v_1 + v_2) + c(v_1 + v_2 + v_3) = 0$. Then $(a + b + c)v_1 + (b + c)v_2 + cv_3 = 0$. By independence, $c = 0$, then $b = 0$, then $a = 0$.

Problem 0.15. For which values of the constants a, b, c, d, e and f are the following vectors linearly independent?

$$\begin{bmatrix} a \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} b \\ c \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} d \\ e \\ f \\ 0 \end{bmatrix}.$$

Solution.

Arrange the three vectors as columns of a 4×3 matrix $M = \begin{bmatrix} a & b & d \\ 0 & c & e \\ 0 & 0 & f \\ 0 & 0 & 0 \end{bmatrix}$. These columns

are independent iff $\text{rank}(M) = 3$, i.e., iff all diagonal entries a, c, f are nonzero. Thus the set is linearly independent exactly when $a \neq 0, c \neq 0$, and $f \neq 0$ (no conditions on b, d, e).

Problem 0.16. Suppose that you are given six vectors $v_1, \dots, v_6 \in \mathbb{R}^5$ with the following properties.

- Any five of them span \mathbb{R}^5 .
- The only relation among them is $v_1 + v_2 + \dots + v_6 = 0$.

Let A be the 5×6 matrix whose column vectors are v_i .

- (a) What is $\dim(\text{Im}(A))$ and $\dim(\ker(A))$?
- (b) Pick any index j . Consider the 5×5 matrix A^j obtained by deleting column j from A . Is A^j invertible?
- (c) Define $T : \mathbb{R}^6 \rightarrow \mathbb{R}^5$ by $T(e_i) = v_i$. For how many distinct scalars c does there exist a linear map $f : \mathbb{R}^5 \rightarrow \mathbb{R}$ with $f(v_i) = c$ for all i ?

Solution.

(a) Any five span $\mathbb{R}^5 \Rightarrow$ the image has dimension 5. By Rank–Nullity on $T : \mathbb{R}^6 \rightarrow \mathbb{R}^5$, $\dim \ker T = 6 - 5 = 1$.

(b) Deleting any one column removes the sole relation $v_1 + \dots + v_6 = 0$, so the remaining 5 columns are independent in \mathbb{R}^5 ; hence each A_j is invertible.

(c) The condition “ $f(v_i) = c$ for all i ” means f is constant on the image of T . Since $\text{Im } T = \mathbb{R}^5$, we must have $f(x) \equiv c$ as a linear map, which forces $c = 0$. Therefore there is exactly one scalar c (namely 0) for which such an f exists.

Problem 0.17. Let $n \geq 2$. Suppose that a linear map $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies $S^2 = 0$ and $\dim(\text{Im}(S)) = k$ for some $k \geq 1$. Define

$$T(\vec{x}) = \vec{x} + S(\vec{x}).$$

- (a) Compute $\dim(\ker T)$ and $\dim(\text{Im } T)$.
- (b) Describe a basis of \mathbb{R}^n in which the matrix of T has the simplest possible block form. State that form.
- (c) For which positive integers m does T^m have the same image as T ? For which m does T^m have the same kernel as T ?

Solution.

(a) $\ker T = \{0\}$ and $\text{Im } T = \mathbb{R}^n$. Indeed, $(I + S)x = 0 \Rightarrow Sx = -x \stackrel{S}{\Rightarrow} 0 = -Sx \Rightarrow x = 0$.

(b) Choose a basis $\{x_i, u_i, y_j\}$ with $Sx_i = u_i, Su_i = 0, Sy_j = 0$; then $[T] = \text{diag}(\underbrace{\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \dots, \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}}_{k \text{ copies}}, I_{n-2k})$.

(c) Since $S^2 = 0$, $(I + S)^m = I + mS$. As in (a), $\ker(I + mS) = \{0\}$ and the image is all of \mathbb{R}^n for every $m \geq 1$.